

Access DB# 76586

## SEARCH REQUEST FORM

Scientific and Technical Information Center

Requester's Full Name: Carmel S. Thompson Examiner #: 79244 Date: 9/26/02  
Art Unit: 1774 Phone Number 305 41488 Serial Number: 09/862 449  
Mail Box and Bldg/Room Location: CP3 11/B/28 Results Format Preferred (circle): PAPER DISK E-MAIL

If more than one search is submitted, please prioritize searches in order of need.

\*\*\*\*\*

Please provide a detailed statement of the search topic, and describe as specifically as possible the subject matter to be searched. Include the elected species or structures, keywords, synonyms, acronyms, and registry numbers, and combine with the concept or utility of the invention. Define any terms that may have a special meaning. Give examples or relevant citations, authors, etc, if known. Please attach a copy of the cover sheet, pertinent claims, and abstract.

Title of Invention: Blue electroluminescence compound for an organic EL device + organic EL device  
Inventors (please provide full names): See attached

Earliest Priority Filing Date: 10/17/2000

\*For Sequence Searches Only\* Please include all pertinent information (parent, child, divisional, or issued patent numbers) along with the appropriate serial number.

Please do a search and a CAS search for the formula in claim 1 where Ar is anthracene.

Please do a search and a CAS search for the formula in claim 5 where Ar is anthracene.

Please search organic EL device using spirobifluorene compounds.

Thanks

## STAFF USE ONLY

	Type of Search	Vendors and cost where applicable
Searcher: <u>ET</u>	NA Sequence (#) _____	STN <u>7160.37</u>
Searcher Phone #: _____	AA Sequence (#) _____	Dialog _____
Searcher Location: _____	Structure (#) <u>(2)</u>	Questel/Orbit _____
Date Searcher Picked Up: _____	Bibliographic _____	Dr. Link _____
Date Completed: <u>9-27-02</u>	Litigation _____	Lexis/Nexis _____
Searcher Prep & Review Time: <u>5</u>	Fulltext _____	Sequence Systems _____
Clerical Prep Time: _____	Patent Family _____	WWW/Internet _____
Online Time: <u>60</u>	Other _____	Other (specify) _____

=> file reg

FILE 'REGISTRY' ENTERED AT 22:50:44 ON 27 SEP 2002  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2002 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file  
provided by InfoChem.

STRUCTURE FILE UPDATES: 26 SEP 2002 HIGHEST RN 455874-53-0  
DICTIONARY FILE UPDATES: 26 SEP 2002 HIGHEST RN 455874-53-0

TSCA INFORMATION NOW CURRENT THROUGH MAY 20, 2002

Please note that search-term pricing does apply when  
conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. See HELP  
PROPERTIES for more information. See STNote 27, Searching Properties  
in the CAS Registry File, for complete details:  
<http://www.cas.org/ONLINE/STN/STNOTES/stnotes27.pdf>

=> d his

(FILE 'HOME' ENTERED AT 22:30:16 ON 27 SEP 2002)

FILE 'LREGISTRY' ENTERED AT 22:30:31 ON 27 SEP 2002  
E SPIROBIFLUORENE/CN

FILE 'REGISTRY' ENTERED AT 22:30:46 ON 27 SEP 2002  
E SPIROBIFLUORENE/CN

L1 1 S E3

FILE 'LREGISTRY' ENTERED AT 22:30:57 ON 27 SEP 2002  
STR 159-66-0

L2

FILE 'REGISTRY' ENTERED AT 22:37:27 ON 27 SEP 2002

L3 0 S L2

L4 STR L2

L5 2 S L4

L6 7 S 2 9841.9.1/RID

FILE 'CAOLD' ENTERED AT 22:41:09 ON 27 SEP 2002  
0 S L6

L7

FILE 'ZCAPLUS' ENTERED AT 22:41:16 ON 27 SEP 2002  
5 S L6

L8

FILE 'LREGISTRY' ENTERED AT 22:42:53 ON 27 SEP 2002  
STR

L9

FILE 'REGISTRY' ENTERED AT 22:47:19 ON 27 SEP 2002

L10 0 S L9  
 L11 STR L9  
 L12 0 S L11  
 L13 1 S L11 FUL  
 SAV L13 THO449/A

FILE 'CAOLD' ENTERED AT 22:49:48 ON 27 SEP 2002

L14 0 S L13



FILE 'ZCAPLUS' ENTERED AT 22:49:58 ON 27 SEP 2002

L15 1 S L13

FILE 'REGISTRY' ENTERED AT 22:50:44 ON 27 SEP 2002

=> d l5 que stat

L4 STR



NODE ATTRIBUTES:

DEFAULT MLEVEL IS ATOM

DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED

NUMBER OF NODES IS 50

STEREO ATTRIBUTES: NONE

L5 2 SEA FILE=REGISTRY SSS SAM L4

100.0% PROCESSED 416 ITERATIONS

2 ANSWERS

SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE \*\*COMPLETE\*\*


BATCH \*\*COMPLETE\*\*

PROJECTED ITERATIONS: 7097 TO 9543

PROJECTED ANSWERS: 2 TO 124

=&gt; d l13 que stat

L11 STR



NODE ATTRIBUTES:

DEFAULT MLEVEL IS ATOM

DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED

NUMBER OF NODES IS 11

STEREO ATTRIBUTES: NONE

L13 1 SEA FILE=REGISTRY SSS FUL L11

100.0% PROCESSED 202930 ITERATIONS

1 ANSWERS

SEARCH TIME: 00.00.07

=&gt; file zcaplus

FILE 'ZCAPLUS' ENTERED AT 22:51:34 ON 27 SEP 2002

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December

26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS is strictly prohibited.

FILE COVERS 1907 - 27 Sep 2002 VOL 137 ISS 14  
FILE LAST UPDATED: 26 Sep 2002 (20020926/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.


CAS roles have been modified effective December 16, 2001. Please check your SDI profiles to see if they need to be revised. For information on CAS roles, enter HELP ROLES at an arrow prompt or use the CAS Roles thesaurus (/RL field) in this file.

=> d l15 1 ibib abs hitstr hitrn

L15 ANSWER 1 OF 1 ZCAPLUS COPYRIGHT 2002 ACS  
ACCESSION NUMBER: 2002:313077 ZCAPLUS  
DOCUMENT NUMBER: 136:332517  
TITLE: Blue phosphors for organic electroluminescent devices  
INVENTOR(S): Kim, Sung Han; Yoo, Han Sung; Kwon, Soon Ki;  
Kim, Yun Hi; Sin, Dong Dhul; Lee, Hyun Uk;  
Chung, Hyung Chul  
PATENT ASSIGNEE(S): Samsung Sdi Co., Ltd., S. Korea  
SOURCE: Jpn. Kokai Tokkyo Koho, 9 pp.  
CODEN: JKXXAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2002121547	A2	20020426	JP 2001-154369	20010523
US 2002055013	A1	20020509	US 2001-862449	20010523
PRIORITY APPLN. INFO.:			KR 2000-60968	A 20001017
OTHER SOURCE(S):	MARPAT 136:332517			
GI				

*application*



*Spirobifluorene*

I


AB The phosphors comprise a spirobifluorene deriv. I (Ar = C6-20 aryl, C6-20 aryl having C1-20 alkyl, C6-20 aryl having C1-20 alkoxy; R1,2 = H, C1-20 alkyl, C6-20 aryl having C1-20 alkyl; C6-20 aryl having C1-20 alkoxy).

IT 413627-08-4

(blue phosphors for org. electroluminescent devices)

RN 413627-08-4 ZCAPLUS

CN Silane, (9,10-anthracenediyl-di-4,1-phenylene)bis(triphenyl- (9CI)  
(CA INDEX NAME)




IT 413627-08-4

(blue phosphors for org. electroluminescent devices)

=> d l8 1-6 cbib abs hitstr hitrn

L8 ANSWER 1 OF 5 ZCAPLUS COPYRIGHT 2002 ACS  
 2002:313077 Document No. 136:332517 Blue phosphors for organic  
 electroluminescent devices. Kim, Sung Han; Yoo, Han Sung; Kwon,  
 Soon Ki; Kim, Yun Hi; Sin, Dong Dhul; Lee, Hyun Uk; Chung, Hyung  
 Chul (Samsung Sdi Co., Ltd., S. Korea). Jpn. Kokai Tokkyo Koho JP  
 2002121547 A2 20020426, 9 pp. (Japanese). CODEN: JKXXAF.  
 APPLICATION: JP 2001-154369 20010523. PRIORITY: KR 2000-60968  
 20001017.

GI



I

AB The phosphors comprise a spirobifluorene deriv. I (Ar = C6-20 aryl,  
 C6-20 aryl having C1-20 alkyl, C6-20 aryl having C1-20 alkoxy; R1,2  
 = H, C1-20 alkyl, C6-20 aryl having C1-20 alkyl; C6-20 aryl having  
 C1-20 alkoxy).


IT 393841-79-7

(blue phosphors for org. electroluminescent devices)


RN 393841-79-7 ZCAPLUS

CN 9,9'-Spirobi[9H-fluorene], 2,2'-(9,10-anthracenediyl)bis[2',7'-  
 bis(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 2-A



IT 393841-79-7

(blue phosphors for org. electroluminescent devices)

L8 ANSWER 2 OF 5 ZCAPLUS COPYRIGHT 2002 ACS  
2001:880187 Document No. 136:158038 Novel blue emitting material with



high color purity. Kim, Yun-Hi; Shin, Dong-Cheol; Kim, Sung-Han; Ko, Chang-Hee; Yu, Han-Sung; Chae, Yun-Soo; Kwon, Soon-Ki (Department of Polymer Science & Engineering and Research Institute of Industrial Technology, Gyeongsang National University, Jinju, 660-701, S. Korea). Advanced Materials (Weinheim, Germany), 13(22), 1690-1693 (English) 2001. CODEN: ADVMEW. ISSN: 0935-9648. Publisher: Wiley-VCH Verlag GmbH.

AB The synthesis of the novel luminescent material 9,10-bis[(2",7"-di-t-butyl)9',9"-spirobifluorenyl]anthracene (TBSA), and the fabrication and performance of a pure blue-emitting org. electroluminescent device (OLED), which has a non-doping structure for full color devices with TBSA as the emitting material, were described. The devices had the configuration of indium tin oxide (ITO)/copper phthalocyanine (CuPc)/1,4-bis[(1-naphthylphenyl)amino]biphenyl (a-NPD)/TBSA/tris(8-hydroxyquinoline) aluminum (Alq3)/LiF/Al, where TBSA was used as the emitting layer, CuPc as the hole-injection layer, a-NPD as the hole-transporting layer (HTL), Alq3 as the electron-transporting layer (ETL), and LiF as the electron-injection layer. The low mol. wt., bis(spirobifluorenyl)anthracene presented a stable non-polymeric org. glass with high glass transition temps., usually assocd. with amorphous polymers. High quality amorphous films of this newly synthesized TBSA with high morphol. stability could be prepd. by vapor deposition. With the newly designed non-doped, blue emitting material in a multilayer device structure, it was possible to achieve a luminous efficiency of 1.22 lm/W at a voltage of 7.7 V and brightness of 300 cd/m<sup>2</sup>. The most important result was the achievement of the purest blue emission nearest to the NTSC std. ever reported.

IT 393841-79-7P


(novel blue emitting material with high color purity)

RN 393841-79-7 ZCAPLUS


CN 9,9'-Spirobi[9H-fluorene], 2,2'-(9,10-anthracenediyl)bis[2',7'-bis(1,1-dimethylethyl)-(9CI) (CA INDEX NAME)

Sept. 2001

PAGE 1-A



PAGE 2-A



IT 393841-79-7P

(novel blue emitting material with high color purity)

L8 ANSWER 3 OF 5 ZCAPLUS COPYRIGHT 2002 ACS

1999:655149 Document No. 132:64193 Closely-spaced chelating centers:

synthesis of novel spiro-bridged bis-phenanthrolines and bis-indole derivatives. Wu, Feiyue; Riesgo, Elvira C.; Thummel, Randolph P.; Juris, Alberto; Hissler, Muriel; El-Ghayoury, Abdelkrim; Ziessel, Raymond (Department of Chemistry, University of Houston, Houston, TX, 77204-5641, USA). Tetrahedron Letters, 40(41), 7311-7314 (English) 1999. CODEN: TELEAY. ISSN: 0040-4039. OTHER SOURCES: CASREACT 132:64193. Publisher: Elsevier Science Ltd..

AB The syntheses of novel sol. ditopic 1,10-phenanthroline ligands bearing a central spiro-[5.5]undecane or a spiro-[5.5]bifluorylidene fragment are reported. The synthetic approach is based on a Friedlander condensation between 8-amino-7-quinolinecarboxaldehyde and either 3,9-diketospiro[5.5]undecane or 2,2'-diacetylspiro[5.5]bifluorylidene derivs. Reaction of the latter with phenylhydrazine and subsequent cyclization afforded 2,2'-di-(2"-indolyl)-[5.5]spirobifluorylidene. The photophys. properties of the new compds. are briefly discussed and Ru(II) and Cu(I) complexes were prepd.

IT 253141-17-2P  
(prepn. of spirocyclic bis(phenanthrolines) and bis(indoles))

RN 253141-17-2 ZCAPLUS

CN Ruthenium(5+), (copper)tetrakis(1,10-phenanthroline-.kappa.N1,.kappa.N10)bis[.mu.-[2,2'-(9,9'-spirobi[9H-fluorene]-2,2'-diyl)bis[1,10-phenanthroline-.kappa.N1,.kappa.N10]]]di-, pentakis[hexafluorophosphate(1-)] (9CI) (CA INDEX NAME)


CM 1

CRN 253141-16-1


CMF C146 H88 Cu N16 Ru2

CCI CCS


PAGE 1-A




PAGE 2-A




PAGE 3-A



PAGE 4-A



PAGE 5-A




CM 2

CRN 16919-18-9

CMF F6 P

CCI CCS



IT 253141-17-2P

(prepn. of spirocyclic bis(phenanthrolines) and bis(indoles))

L8 ANSWER 4 OF 5 ZCAPLUS COPYRIGHT 2002 ACS

1997:343394 Document No. 127:127809 Electrochemistry of 9,9'-spirobifluorene derivatives: electrosynthesis of stereoisomeric 2,3-bis(2'-acetyl-9,9'-spirobifluoren-2-yl)butane-2,3-diols and of 1-(2'-acetyl-9,9'-spirobifluoren-2-yl)ethanol and redox properties of polyacetylated spirobifluorenes. Mattiello, Leonardo; Rampazzo, Liliana (Dep. of ICMMPM, Sede Chimica, Universita di Roma "La Sapienza", Rome, 00161, Italy). Electrochimica Acta, 42(13-14), 2257-2264 (English) 1997. CODEN: ELCAAV. ISSN: 0013-4686. Publisher: Elsevier.


AB 2,2'-Diacetyl-9,9'-spirobifluorene 3a, an axially dissym. compd., when dissolved in virtually aprotic solvent DMF, was characterized by several redn. peaks in the voltammetric expts., using a glassy carbon electrode; the 1st two electron transfers are reversible, with std. potentials  $E_{10} = -1.75$  V and  $E_{20} = -1.90$  V vs. SCE resp. A 3rd redn. step occurs with  $E_p = -2.40$  V, at  $\nu. = 50$  mV s<sup>-1</sup>. The 1st two steps are pertinent to the formation of the anion radical A.bul.- and of the dianion diradical -.bul.A-.bul. of 3a (A), resp. Controlled potential electrolysis of 3a on a rvc (reticulated vitreous carbon) electrode in DMF-Et<sub>4</sub>NClO<sub>4</sub> (0.1M), with acetic acid 1:1 mol 3a/mol acid added, furnishes the two diastereomeric keto-alcs. and the six diastereomeric keto-pinacols. This result was obtained if the electrolysis is stopped after 1 F/mol 3a was passed. If the electrolysis is continued until both C=O groups are reduced, a mixt. of diastereomeric pinacols was obtained. Spectroscopic properties of the diastereomeric keto-pinacols are sensitive to the various spatial arrangements in the isomers. Some aspects of the redox properties of 3a, as related to the spiro-structure of the mol., are also discussed. Redox properties of tri- and tetra-acetylated spirobifluorenes in virtually aprotic conditions, as detd. through cyclic voltammetry, are in turn related to those of the diacetylated deriv. 3a: 2,2'-diacetyl-9,9'-spirobifluorene and the tri- and tetra-acetylated spirobifluorenes are the first examples of spiro compds. bearing multiple redox-active substituents.

IT 192565-89-2

(stereoisomers; prepn. by controlled potential electrolysis of diacetylspirobifluorene on glassy carbon in DMF contg. acetic acid)

RN 192565-89-2 ZCAPLUS

CN Ethanone, 1,1'-[(1,2-dihydroxy-1,2-dimethyl-1,2-ethanediyl)bis(9,9'-spirobi[9H-fluorene]-2',2-diyl)]bis- (9CI) (CA INDEX NAME)




IT 192565-90-5P

(stereoisomers; prepn. by electrochem. redn. of diacetylspirobifluorene on glassy carbon in DMF contg. acetic acid)

RN 192565-90-5 ZCAPLUS

CN 9,9'-Spirobi[9H-fluorene]-2,2'-dimethanol, .alpha.-[1-hydroxy-1-[2'-(1-hydroxyethyl)-9,9'-spirobi[9H-fluoren]-2-yl]ethyl]-.alpha.,.alpha.'-dimethyl- (9CI) (CA INDEX NAME)



IT 192565-89-2

(stereoisomers; prepn. by controlled potential electrolysis of diacetylspirobifluorene on glassy carbon in DMF contg. acetic acid)


IT 192565-90-5P

(stereoisomers; prepn. by electrochem. redn. of diacetylspirobifluorene on glassy carbon in DMF contg. acetic acid)

L8 ANSWER 5 OF 5 ZCAPLUS COPYRIGHT 2002 ACS

1994:163140 Document No. 120:163140 Electrochemistry of 9,9'-spirobifluorene derivatives: 2-acetyl- and 2,2'-diacetyl-9,9'-spirobifluorene. Preparation of stereoisomeric 2,3-bis(9,9'-spirobi-fluoren-2-yl)butane-2,3-diols. Mattiello, Leonardo; Rampazzo, Liliana (Univ. Roma 'La Sapienza', Rome, 00161, Italy). Journal of the Chemical Society, Perkin Transactions 2: Physical Organic Chemistry (1972-1999) (11), 2243-7 (English) 1993. CODEN: JCPKBH. ISSN: 0300-9580. OTHER SOURCES: CASREACT 120:163140.

GI



AB 2-Acetyl- and 2,2'-diacetyl-9,9'-spirobifluorene 1 and 2 were studied by cyclic voltammetry in DMF. The corresponding anion radicals show remarkable persistency in aprotic DMF. The (apparent) std. potentials are E.degree. = -1.77 V (SCE) and E.degree. = -1.75 V for the (quasi-reversible) redn. of 1 and 2 to the anion radicals, resp. Preparative electrolysis of 1 in DMF-Et<sub>4</sub>NClO<sub>4</sub> (0.1 mol dm<sup>-3</sup>), with excess acetic acid as proton donor, furnished alc. I and the diastereoisomeric pinacols [meso- and (.+-.)-II], which were isolated and characterized. The diastereoisomeric excess, de, as evaluated (NMR) or the electrolyzed soln. was only slightly in favor of the (.+-.)- compd. Spectroscopic properties of compds. 1,2, I-II are, inter alia, the <sup>13</sup>C NMR chem. shift for the spiro-carbon at .delta. = 65.9 (TMS), and the fragmentation patterns in the mass spectra, with the 100% relative abundance of the mol.-ion M.bul.+ in the case of the arom. ketones 1 and 2. Some comments on the influence of conformations of meso- and (.+-.)-II and the presence of an intramol. hydrogen-bond. in some intermediates and products are also presented.


IT 153335-20-7P 153335-21-8P  
(prepn. of)

RN 153335-20-7 ZCAPLUS

CN 2,3-Butanediol, 2,3-bis(9,9'-spirobi[9H-fluoren]-2-yl)-, (R\*,S\*)-(9CI) (CA INDEX NAME)

Relative stereochemistry.






RN 153335-21-8 ZCAPLUS

CN 2,3-Butanediol, 2,3-bis(9,9'-spirobi[9H-fluoren]-2-yl)-, (R\*,R\*)-  
(9CI) (CA INDEX NAME)

Relative stereochemistry.



IT 153335-20-7P 153335-21-8P  
(prepn. of)